

POULTRY FEEDS GUIDE







Country Junction produces feed for both major producers and smaller operations, with specialization in making premixed, customized feeding programs for all markets, including organic poultry producers.

OUR FEEDS CONTAIN A PROPER BALANCE OF NUTRIENTS FOR EACH LIFE STAGE OF POULTRY PRODUCTION, WITH RATIONS FOR:

- Starting, growing and finishing meat-type birds
- Starting and developing pullets to becoming laying hens
- Laying hens to increase egg-production and shell quality

FEATURES AND BENEFITS

Though each feed contains its individual advantages you can be sure that from Country Junction your feed will be

- Nutritionally Balanced
- Formulated to provide the protein, energy, vitamins, and minerals in an optimal balance specific to the feeds purpose
- Our poultry and fowl feeds are formulated with whole grains and vegetable proteins without the use of animal byproducts
- The grains and grain products used in these feeds are from our Canadian Prairie region

POULTRY





Crude Protein (Min.)	20.0%	Sodium (Actual)	0.16%
Crude Fat (Min.)	2.0%	Vitamin A (Min.)	10,500 IU/kg
Crude Fiber (Max.)	6.0%	Vitamin D3 (Min.)	2,500 IU/kg
Calcium (Actual)	1.00%	Vitamin E (Min.)	35 IU/kg
Phosphorus (Actual)			

20% CHICK STARTER CRUMBLE

FEED FORM: Crumble **PRODUCT #:** P100000B *This feed contains added selenium at 0.3 mg/kg.

FEEDING DIRECTIONS

20% Chick Starter should be fed as the sole ration to broiler chicks up to 3 weeks of age or to replacement pullets up until 6 weeks of age. Oyster shell or insoluble grit could be sprinkled on the feed once a week at a rate of 0.5 to 1.0 kg per 100 birds. Allow chicks access to fresh water at all times. Replacement layer chicks will eat about 1 kg of starter in 6 weeks. Broiler chicks will eat about 1 kg of starter in 3 weeks.

- Non-medicated feed
- Dual purpose starter
 - Can be used to start laying pullets as well as meat birds
- Crumbled Feed



16% POULTRY GROWER CRACKED WHEAT

FEED FORM: Mash PRODUCT #: P612100B *This feed contains added selenium at 0.3 mg/kg.

GUARANTEED ANALYSIS

Crude Protein (Min.)	16.0%	Sodium (Actual)	0.16%
Crude Fat (Min.)	2.5%	Vitamin A (Min.)	12,000 IU/kg
Crude Fiber (Max.)	6.5%	Vitamin D3 (Min.)	2,900 IU/kg
Calcium (Actual)	1.0%	Vitamin E (Min.)	40 IU/kg
Phosphorus (Actual)	0.6%		

FEEDING DIRECTIONS

Country Junction 16% Poultry Grower should be fed as the sole ration to replacement laying chicks from 7 to 18 weeks of age. This 16% Poultry Grower may be fed to broiler (Meat Type) growing chickens from 3 weeks of age to market weight. Oyster shell or insoluble grit could be sprinkled on the feed once a week at the rate of 0.5 to 1.0 kg per 100 birds. Allow birds access to fresh water at all times.

- Non-medicated feed
- Dual purpose grower
 - Can be used to grow laying pullets as well as finishing your meat birds
- Contains cracked wheat and rolled barley



Crude Protein (Min.)	17.0%	Sodium (Actual)	0.16%
Crude Fat (Min.)	2.0%	Vitamin A (Min.)	18,500 IU/kg
Crude Fiber (Max.)	4.5%	Vitamin D3 (Min.)	4,000 IU/kg
Calcium (Actual)	3.7%	Vitamin E (Min.)	70 IU/kg
Phosphorus (Actual)	0.6%		

17% LAYER RATION CRACKED WHEAT

FEED FORM: Mash PRODUCT #: P632110B *This feed contains added selenium at 0.3 mg/kg.

FEEDING DIRECTIONS

17% Layer Diet is designed to be fed to laying hens from approximately 5% lay to end of lay. This diet is intended to be fed at approximately 95 - 110 grams per hen per day. The exact feeding rate will be determined by the strain of bird, body weight and condition and barn temperature. Oyster shell or insoluble grit could be sprinkled on the feed once a week at the rate of 0.5 to 1.0 kg per 100 birds. Allow hens access to fresh water at all times.

- Non-medicated feed
- Good for Multi-aged flocks
- Contains cracked wheat



LAYER SUPPLEMENT

FEED FORM: Crumble **PRODUCT #:** L200100 *This feed contains added selenium at 1.2 mg/kg.

GUARANTEED ANALYSIS

Crude Protein (Min.)	28.0%	Copper (Actual)	80 mg/kg
Crude Fat (Min.)	3.0%	Iodine (Actual)	10 mg/kg
Crude Fiber (Max.)	6.0%	Iron (Actual)	260 mg/kg
Calcium (Actual)	15.2%	Manganese (Actual)	520 mg/kg
Phosphorus (Actual)	1.4%	Zinc (Actual)	440 mg/kg
Phosphorus (Actual) Potassium (Actual)	1.4% 0.6%	Zinc (Actual) Vitamin A (Min.)	440 mg/kg 35,500 IU/kg
			0. 0

FEEDING DIRECTIONS

Feed the following suggested rations as the sole ration. Country Junction Layer Supplement can be mixed with farm grains to produce a layer ration for non-commercial laying hens.

Total	1000 kg	1000 kg
Layer Supplement	250 kg	250 kg
Barley		300 kg
Wheat	750 kg	450 kg
	Layer Diet I (18 - 30 weeks)	Layer Diet II (30 - 52 weeks)

Layer Diet I will be used at the start of the laying cycle, which can be replaced with Layer Diet II after the birds have been laying for 3 to 4 months, depending upon type of bird, weight and number of eggs collected. Always allow birds access to clean water and free choice poultry grit or oystershell at all times.

- Non-medicated supplement
- Good for Multi-aged flocks
- Can be mixed with farm grains to produce a layer ration for non-commercial laying hens





TURKEY STARTER

FEED FORM: Crumble PRODUCT #: T100008B *This feed contains added selenium at 0.3 mg/kg.

GUARANTEED ANALYSIS

Crude Protein (Min.)	26.0%	Sodium (Actual)	0.16%
Crude Fat (Min.)	3.0%	Vitamin A (Min.)	15,000 IU/kg
Crude Fiber (Max.)	8.0%	Vitamin D3 (Min.)	4,500 IU/kg
Calcium (Actual)	1.38%	Vitamin E (Min.)	50 IU/kg
Phosphorus (Actual)	0.95%		

FEEDING DIRECTIONS

Feed Country Junction Turkey Starter to turkey poults from arrival to 6 or 8 weeks of age. May also be fed to ducks, geese, and game birds up to 4 weeks of age. Oyster shell or insoluble grit could be sprinkled on the feed once a week at a rate of 0.5 to 1.0kg/100 birds. Ensure the poults are warm, dry, have adequate fresh air and free choice access to clean water.

• Turkey hen poults will eat 4-5 kg of starter in 7 weeks

• Turkey tom poults will eat 5-6 kg of starter in 7 weeks

- High protein starter
- Crumbled feed



Crude Protein (Min.)	22.0%	Sodium (Actual)	0.17%
Crude Fat (Min.)	3.0%	Vitamin A (Min.)	15,000 IU/kg
Crude Fiber (Max.)	7.0%	Vitamin D3 (Min.)	3,800 IU/kg
Calcium (Actual)	1.2%	Vitamin E (Min.)	50 IU/kg
Phosphorus (Actual)	0.82%		

TURKEY GROWER

 FEED FORM:
 Pellet
 PRODUCT #: T111000B

 *This feed contains added selenium at 0.3 mg/kg.

FEEDING DIRECTIONS

Feed Country Junction Turkey Grower to turkey poults from 42 days of age to 70 days of age or to market weight. Ensure the poults are warm, dry, have adequate fresh air and free choice access to clean water. After 70 days of age switch, to Country Junction Turkey Finisher.

FEATURES AND BENEFITS

• Non-medicated

Pelleted feed



DUCK & GOOSE



Crude Protein (Min.)	21.0%	Sodium (Actual)	0.18%
Crude Fat (Min.)	3.0%	Vitamin A (Min.)	15,000 IU/kg
Crude Fiber (Max.)	5.0%	Vitamin D3 (Min.)	3,600 IU/kg
Calcium (Actual)	0.8%	Vitamin E (Min.)	35 IU/kg
Phosphorus (Actual)	0.65%		

21% DUCK AND GOOSE STARTER

FEED FORM: Crumble **PRODUCT #:** P400000B *This feed contains added selenium at 0.3 mg/kg.

FEEDING DIRECTIONS

Country Junction Duck & Goose Starter should be fed as the sole ration to ducklings and goslings up until 4 weeks of age. Oyster shell or insoluble grit could be sprinkled on the feed once a week at the rate of 0.5 to 1.0 kg per 100 birds. Allow ducklings and goslings access to fresh water at all times.

Ducklings will eat about 3-3.6 kg of starter in 4 weeks depending upon the breed and sex of the birds.

Goslings will eat about 4-5 kg of starter in 4 weeks depending upon breed and sex of the birds.

- Non-medicated feed
- Dual purpose starter which can be used to start meat ducks and geese or young waterfowl that will be laying and/or breeding
- Crumbled Feed



DUCK & GOOSE GROWER

FEED FORM: Pellet **PRODUCT #:** P401000B *This feed contains added selenium at 0.3 mg/kg.



GUARANTEED ANALYSIS

Crude Protein (Min.)	18.0%	Sodium (Actual)	0.16%
Crude Fat (Min.)	2.0%	Vitamin A (Min.)	13,500 IU/kg
Crude Fiber (Max.)	5.0%	Vitamin D3 (Min.)	2,900 IU/kg
Calcium (Actual)	0.9%	Vitamin E (Min.)	45 IU/kg
Phosphorus (Actual)	0.6%		

FEEDING DIRECTIONS

Country Junction Duck & Goose Grower should be fed as the sole ration to ducklings and goslings from 5 weeks of age to market weight. Oyster shell or insoluble grit could be sprinkled on the feed once a week at a rate of 0.5 to 1.0 kg per 100 birds. Allow ducklings and goslings access to fresh water at all times.

- Non-medicated
- Pelleted Feed



For a list of other bagged poultry & fowl feeds available in 2 tonne minimums, as well as a large variety of other bagged feeds please contact your Country Junction Feeds representative.

Also available cracked chicken scratch by the bag.



4707 40 Ave , Wetaskiwin, AB T9A 2B8 Phone: +1 800-729-9155 Mill: 780-352-9155 Email: info@countryjunctionfeeds.com